The Bryologist 90(4), 1987, pp. 468–474
Copyright © 1987 by the American Bryological and Lichenological Society, Inc.

Index to Volume 90 of THE BRYOLOGIST[1,2]

Abies lasiocarpa, epiphytic lichens on+, 212
abscisic acid, effect on stomata of Anthoceros spp.+, 393; in Anthocerotae and Marchantiales, 393+
Acanthocoleus aberrans, 339+; juddii, 339+
Acarospora nodulosa var. nodulosa, new to North America, 48
acid precipitation, effects on Sphagnum species, 221
Acolea lacerata, 362
Acrobolbus ciliatus, 362, mapped, 363
Acrocordia ambigua (syn.), *163*; **megalospora**, comb. nov., 163
Acrolejeunea emergens, 339+; heterophylla, 341; torulosa, 339+
AKIYAMA, H. Morphology and anatomy of the peristome in Dozya japonica (Musci) and reconsideration of its affinity to the Leucodontaceae, 409
Alabama, Syrrhopodon prolifer var. papillosus new to, 263; Syrrhopodon species in, 263+
algae, associates in Catapyrenium species, 27; associates in Dermatocarpon species, 27; associates in Placidiopsis species, 38; blue-green, response to desiccation, 241+; green, response to desiccation, 241+
ALLEN, B. H. (review). Noel H. Holmgren & Bobbi Angell, Botanical Illustrations: Preparation for Publication, 198
ALLEN, B. H. Hookeriopsis **crosbyi** (Musci: Hookeriaceae): a new species from western Panama, 45
amber, fossil of Stictolejeunea cf. squamata, 339
ANDERSON, L. E. A new species of Polytrichum from the southern Appalachians, 1
Andreaea alpina, 23; blyttii, 25; crassinervia, 19+; frigida, 15+; heinemannii, 17+; **megistospora**, sp. nov., 18; **megistospora** ssp. **epapillosa**, 20*, 21*, ssp. nov., 23, ecology, 24, mapped, 24; megistospora ssp. megistospora, 18+, 20*, 21*, ecology, 22, mapped, 23; nivalis, 17; rothii ssp. falcata, 19+; rothii ssp. huntii, 25; rothii ssp. rothii, 17+; rupestris var. rupestris, 23; **schofieldiana**, sp. nov., 15+, 16*, ecology and distribution of, 17, mapped, 18; sinuosa, 22
Andreaeae, key to European, northern North American costate species of, 25; sect. Nerviae, 15+, spore size in systematics of, 25
ANDRUS, R. E. Nomenclatural changes in Sphagnum imbricatum sensu lato, 217
Anisomeridium **albisedum**, comb. nov., 163; **ambiguum**, comb. nov., 163; **carinthiacum**, comb. nov., 163; **distans**, comb. nov., 163; **leucochlorum**, comb. nov., 163; **sanfordense**, comb. nov., 163
Anthoceros crispulus, 394; dichotomus, 393+; laevis, 393+; punctatus, 393+; abscisic acid in, 393+
Anthracothecium eschweileri (syn.), 163
Anthracothecium **prasinum**, comb. nov., 163
apical organization, in Haplomitrium mnioides, 383
Appalachian Mountains, Bartleya expression of Dicranella cerviculata in, 43+; peatlands, acid precipitation a threat to, 221; Polytrichum **appalachianum**, sp. nov. in, 1+
Aptychus chlorocormus, 248
arabinogalactan proteins, in Hepaticae, 401
Arachniopsis coactilis, 374
Archilejeunea auberiana, 341; crispistipula, 339+; fuscescens, 339+; juliformis (syn.), *339*; parviflora, 341; porelloides, 339+
Arthopyrenia ambigua (syn.), *163*; carinthiaca (syn.), *163*; leucochlora (syn.), *163*; pleiomerella (syn.), *164*; **plumbaria**, comb. nov., 163; sanfordensis (syn.), *163*; submuriformis (syn.), *164*
Ascidiota blepharophylla ssp. alaskana, 345+
Aspicilia **annulata**, comb. nov., 163; **caesiopruinosa**, comb. nov., 163; **limitata**, comb. nov., 163; **mastoidea**, comb. nov., 163; **narssaquensis**, comb. nov., 163; **pertusa**, comb. nov., 163
Asterella californica, 345+; palmeri, 345+
Astomiopsis, a revision of, 309; amblyocalyx, 309+, 312*; exserta, 309+; **exserta**, comb. nov., 316, 317*; key to species of, 311; kieneri (syn.), *309*+; pacifica, 309+, 314*; sinensis, 309+, 319*; subulata, 315*
Astrella angusta, 251; blumeana, 251
Atrichum anomalum, 3
AUSTIN, K. A. & R. K. WIEDER. Effects of elevated H^+, SO_4^{2-}, NO_3^-, and NH_4^+ in simulated acid precipitation on the growth and chlorophyll content of 3 North American Sphagnum species, 221
Australia, Lord Howe Island, Frullania species of, 365; Monoclea species in, 371+
auxin, in phototropism of Pellia epiphylla, 390+

Barbula afro-duriuscula (syn.), *234*
BARTHOLOMEW, S. E. Apical organization and leaf ontogeny in Haplomitrium mnioides (Lindb.) Schust., 383
Bartleya expression of Dicranella cerviculata, distribution of, 44
Bartleya ohioensis, 43+
BASILE, D. V. & M. R. BASILE. The occurrence of cell wall-associated arabinogalactan proteins in the Hepaticae, 401
BASILE, M. R. *See* BASILE, D. V. & M. R. BASILE, 401
Bazzania ambigua, 345+
Bestia, relationships of, 40+; longipes, 40+, placed in Brachytheciaceae, 40; vancouveriensis, 40+, placed in Thamnobryaceae, 40
Biatora endocyanea (syn.), *164*
biogeography, of transoceanic species of Lejeuneaceae subfam. Ptychanthoideae, 337
BISANG, I. & R. SCHUMACKER. Southwesternmost North American locality of Odontoschisma macounii (Aust.) Underw. and Scapania spitzbergensis (Lindb.) K. Mull. at Sutherland Falls (Blanket Creek Provincial Park, British Columbia, Canada), 237
BISCHLER, H. The typification and synonymy of Japanese Marchantias described by Stephani, 331
Blanket Creek Provincial Park, British Columbia, rare liverworts in, 237+
Blasia pusilla, spermatogenesis in, 419+
Blepharolejeunea chimantaensis, 338+; fuegiana, 341; incongrua, 339+; securifolia, 339
blepharoplast assembly, in bryophyte spermatogenesis, 439+
Blepharostoma arachnoideum, 345+, mapped, 346
Bottaria composita (syn.), *164*; cruentata (syn.), *164*
Brachiolejeunea laxifolia, 339; phyllorhiza, 339+, mapped, 340

[1] Compiled by P. M. Eckel, Clinton Herbarium, Buffalo Museum of Science, Buffalo, NY 14211.

[2] New scientific names and new combinations are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign; italicized page numbers indicate taxa in synonymy.

0007-2745/87/000–000$0.00/0

The Bryologist 90(4), 1987, pp. 468–474
Copyright © 1987 by the American Bryological and Lichenological Society, Inc.

Index to Volume 90 of THE BRYOLOGIST[1,2]

Abies lasiocarpa, epiphytic lichens on+, 212
abscisic acid, effect on stomata of Anthoceros spp.+, 393; in Anthocerotae and Marchantiales, 393+
Acanthocoleus aberrans, 339+; juddii, 339+
Acarospora nodulosa var. nodulosa, new to North America, 48
acid precipitation, effects on Sphagnum species, 221
Acolea lacerata, 362
Acrobolbus ciliatus, 362, mapped, 363
Acrocordia ambigua (syn.), *163*; **megalospora**, comb. nov., 163
Acrolejeunea emergens, 339+; heterophylla, 341; torulosa, 339+
AKIYAMA, H. Morphology and anatomy of the peristome in Dozya japonica (Musci) and reconsideration of its affinity to the Leucodontaceae, 409
Alabama, Syrrhopodon prolifer var. papillosus new to, 263; Syrrhopodon species in, 263+
algae, associates in Catapyrenium species, 27; associates in Dermatocarpon species, 27; associates in Placidiopsis species, 38; blue-green, response to desiccation, 241+; green, response to desiccation, 241+
ALLEN, B. H. (review). Noel H. Holmgren & Bobbi Angell, Botanical Illustrations: Preparation for Publication, 198
ALLEN, B. H. Hookeriopsis **crosbyi** (Musci: Hookeriaceae): a new species from western Panama, 45
amber, fossil of Stictolejeunea cf. squamata, 339
ANDERSON, L. E. A new species of Polytrichum from the southern Appalachians, 1
Andreaea alpina, 23; blyttii, 25; crassinervia, 19+; frigida, 15+; heinemannii, 17+; **megistospora**, sp. nov., 18; **megistospora** ssp. **epapillosa**, 20*, 21*, ssp. nov., 23, ecology, 24, mapped, 24; megistospora ssp. megistospora, 18+, 20*, 21*, ecology, 22, mapped, 23; nivalis, 17; rothii ssp. falcata, 19+; rothii ssp. huntii, 25; rothii ssp. rothii, 17+; rupestris var. rupestris, 23; **schofieldiana**, sp. nov., 15+, 16*, ecology and distribution of, 17, mapped, 18; sinuosa, 22
Andreaeae, key to European, northern North American costate species of, 25; sect. Nerviae, 15+, spore size in systematics of, 25
ANDRUS, R. E. Nomenclatural changes in Sphagnum imbricatum sensu lato, 217
Anisomeridium **albisedum**, comb. nov., 163; **ambiguum**, comb. nov., 163; **carinthiacum**, comb. nov., 163; **distans**, comb. nov., 163; **leucochlorum**, comb. nov., 163; **sanfordense**, comb. nov., 163
Anthoceros crispulus, 394; dichotomus, 393+; laevis, 393+; punctatus, 393+; abscisic acid in, 393+
Anthracothecium eschweileri (syn.), 163
Anthracothecium **prasinum**, comb. nov., 163
apical organization, in Haplomitrium mnioides, 383
Appalachian Mountains, Bartleya expression of Dicranella cerviculata in, 43+; peatlands, acid precipitation a threat to, 221; Polytrichum **appalachianum**, sp. nov. in, 1+
Aptychus chlorocormus, 248
arabinogalactan proteins, in Hepaticae, 401
Arachniopsis coactilis, 374
Archilejeunea auberiana, 341; crispistipula, 339+; fuscescens, 339+; juliformis (syn.), *339*; parviflora, 341; porelloides, 339+
Arthopyrenia ambigua (syn.), *163*; carinthiaca (syn.), *163*; leucochlora (syn.), *163*; pleiomerella (syn.), *164*; **plumbaria**, comb. nov., 163; sanfordensis (syn.), *163*; submuriformis (syn.), *164*
Ascidiota blepharophylla ssp. alaskana, 345+
Aspicilia **annulata**, comb. nov., 163; **caesiopruinosa**, comb. nov., 163; **limitata**, comb. nov., 163; **mastoidea**, comb. nov., 163; **narssaquensis**, comb. nov., 163; **pertusa**, comb. nov., 163
Asterella californica, 345+; palmeri, 345+
Astomiopsis, a revision of, 309; amblyocalyx, 309+, 312*; exserta, 309+; **exserta**, comb. nov., 316, 317*; key to species of, 311; kieneri (syn.), *309*+; pacifica, 309+, 314*; sinensis, 309+, 319*; subulata, 315*
Astrella angusta, 251; blumeana, 251
Atrichum anomalum, 3
AUSTIN, K. A. & R. K. WIEDER. Effects of elevated H^+, SO_4^{2-}, NO_3^-, and NH_4^+ in simulated acid precipitation on the growth and chlorophyll content of 3 North American Sphagnum species, 221
Australia, Lord Howe Island, Frullania species of, 365; Monoclea species in, 371+
auxin, in phototropism of Pellia epiphylla, 390+

Barbula afro-duriuscula (syn.), *234*
BARTHOLOMEW, S. E. Apical organization and leaf ontogeny in Haplomitrium mnioides (Lindb.) Schust., 383
Bartleya expression of Dicranella cerviculata, distribution of, 44
Bartleya ohioensis, 43+
BASILE, D. V. & M. R. BASILE. The occurrence of cell wall-associated arabinogalactan proteins in the Hepaticae, 401
BASILE, M. R. *See* BASILE, D. V. & M. R. BASILE, 401
Bazzania ambigua, 345+
Bestia, relationships of, 40+; longipes, 40+, placed in Brachytheciaceae, 40; vancouveriensis, 40+, placed in Thamnobryaceae, 40
Biatora endocyanea (syn.), *164*
biogeography, of transoceanic species of Lejeuneaceae subfam. Ptychanthoideae, 337
BISANG, I. & R. SCHUMACKER. Southwesternmost North American locality of Odontoschisma macounii (Aust.) Underw. and Scapania spitzbergensis (Lindb.) K. Mull. at Sutherland Falls (Blanket Creek Provincial Park, British Columbia, Canada), 237
BISCHLER, H. The typification and synonymy of Japanese Marchantias described by Stephani, 331
Blanket Creek Provincial Park, British Columbia, rare liverworts in, 237+
Blasia pusilla, spermatogenesis in, 419+
Blepharolejeunea chimantaensis, 338+; fuegiana, 341; incongrua, 339+; securifolia, 339
blepharoplast assembly, in bryophyte spermatogenesis, 439+
Blepharostoma arachnoideum, 345+, mapped, 346
Bottaria composita (syn.), *164*; cruentata (syn.), *164*
Brachiolejeunea laxifolia, 339; phyllorhiza, 339+, mapped, 340

[1] Compiled by P. M. Eckel, Clinton Herbarium, Buffalo Museum of Science, Buffalo, NY 14211.

[2] New scientific names and new combinations are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign; italicized page numbers indicate taxa in synonymy.

0007-2745/87/000–000$0.00/0

Brachymitrion moritzianum, 247
Brazil, Zoopsis antillana, new to, 374+; new state records in, 374+
British Columbia, rare hepatics in Blanket Creek Provincial Park, Sutherland Falls, 237+
Bryomanginia, 309+
Bryopteris diffusa, 374+; fruticulosa, 375
Bryoria lanestris, 212+
Bulbothrix confoederata, 417; coronata, 417; decurtata, 417; goebelii, 417; isidiza, distribution in the United States, 417+; laevigatula, 417; tabacina, 417

Callicosta bipinnata, 247; fendleri, 249
Calobryum blumei, 383; gibbsiae, 383
Calymperaceae, Pottiaceae and Serpotortellaceae compared with, 232
Calymperes, new synonyms and range records for paleotropic species in, 201+; acanthoneuron (syn.), *205*; afzelii, new synonyms of, 203+; annamense (syn.), *203*; ascensionis (syn.), *206*; beccarii, 205; bolomboense (syn.), *206*; borgenii (syn.), *208*; borgenii var. congolense (syn.), *205*; burmense (syn.), *206*; cacazouense (syn.), *208*; casamancae (syn.), *206*; caudatum (syn.), *207*; ceylanicum (syn.), *207*; chevalieri (syn.), *205*; chevalieri var. angustifolium (syn.), *205*; cochlearifolium (syn.), *206*; congolense (syn.), *205*; corbieri (syn.), *206*; cucullatum (syn.), *206*; decolorans, new synonyms of, 205; decolorans var. latifolium (syn.), *205*; derelictum (syn.), *208*; dilatatum (syn.), *205*; erosulum (syn.), *205*; erosum, new synonyms of, 206; exlimbatum (syn.), *209*; gabonense (syn.), *206*; gilletii (syn.), *206*; gilletii var. angustifolium (syn.), *203*; guildingii (syn.), *247*; heterophyllum (syn.), *206*; hildebrandtii (syn.), *203*; hispidum (syn.), *208*; hyalinolimbatum (syn.), *209*; integrifolium (syn.), *203*; lamellosulum (syn.), *209*; leucomitrium (syn.), *209*; ligulare (syn.), *207*; linealifolium (syn.), *209*; lonchophyllum, new synonyms of, 206; loucoubense var. minus (syn.), *205*; mahense (syn.), *205*; malimbae (syn.), *206*; maroant setrae (syn.), *203*; maroantsetrae var. angustifolium (syn.), *206*; mathieui (syn.), *206*; mayottense (syn.), *209*; microdictyon (syn.), *203*; nossibeanum (syn.), *207*; obtusatum (syn.), *205*; occidentale (syn.), *209*; omanicum (syn.), *209*; pachyphyllum (syn.), *205*; palisottii, new synonyms of, 206; palmicola (syn.), *209*; perlimbatum (syn.), *203*; perrottetii (syn.), *203*; perserratum (syn.), *206*; pobeguinii (syn.), *205*; polii (syn.), *209*; porrectum, 207; principis (syn.), *207*; pygmaeum (syn.), *209*; quintasii (syn.), *203*; rabenhorstii (syn.), *205*; reductum (syn.), *209*; retusum (syn.), *201*; rhodesiae (syn.), *203*; robillardii (syn.), *209*; robinsonii, range extension of, 208; sanctaemariae (syn.), *207*; sarawakensis (syn.), *203*; schmidtii, 207; seychellarum (syn.), *209*+; sikkimense (syn.), *208*; stenophyllum (syn.), *207*; subcrassilimbatum (syn.), *206*; subdecolorans (syn.), *205*; subdiaphanum (syn.), *207*; taitense, new synonyms of, 207; tenerrimum (syn.), *210*; tenerum, 207+, new synonyms of, 207; usambaricum (syn.), *203*; victoriae (syn.), *203*; zanzibarense (syn.), *207*
Calyptothecium duplicatum, 247
Campbell, E. O. Monoclea (Hepaticae); distribution and number of species, 371
Campylopus concolor, 247; densicoma, 247; fragilis, 247; richardii, 249; tallulensis, 247
Caron, P. J. *See* Thomas, R. J., P. J. Caron & R. S. Watt, 390
Catapyrenium, in North America, 27+; key to the North American species of, 28; **acarosporoides**, 27+, comb. nov., 28, 29*, mapped, 31; **caeruleopulvinum**, 27+, 29*, sp. nov., 30, mapped, 31; cinereum, 27+, 29*, mapped, 32; compactum, 27+, 29*, mapped, 33; **granulosum**, 27+, 29*, comb. nov., 33, mapped, 31; **heppioides**, 27+, 29*, comb. nov., 33; lachneum, 27+, 30*, mapped, 34; michelii, 28+, 30*, mapped, 35; **plumbeum**, 27+, 30*, comb. nov., 36, mapped, 36; **squamellum**, 27+, 30*, comb. nov., 36; **tuckermanii**, 27+, 30*, comb. nov., 36, mapped, 36; **zahlbruckneri**, 27+, 30*, comb. nov., 38, mapped, 38
Catinaria **subcorallina**, comb. nov., 163
Caudalejeunea lehmanniana, 339
cell walls, arabinogalactan proteins in hepatic, 401
Cephalozia brinkmanii, mapped, 346
Cephaloziella brinkmanii, 345+
Cetraria aurescens (syn.), *164*; juniperina var. canadensis (syn.), *164*; merrillii (syn.), *164*; oakesiana (syn.), *164*; pallidula (syn.), *164*; pinastri (syn.), *213*+; platyphylla (syn.), *164*; viridis (syn.), *164*
chromatin condensation, developmental aspects in liverworts, 376
Churchill, S. P. & I. Sastre-De Jesús. Nuevos registros de musgos para los departamentos de Antioquia y Choco, Colombia, y una nueva especie del genero Trichosteleum, 246
coal deposits, Dicranella cerviculata-form associated with, 43+
Coal Moss expression of Dicranella cerviculata, distribution of, 44
Cololejeunea macounii, 344+
Colombia, new records to Dept. of Antioquia, 246; new records to Dept. of Choco, 246; species new to, 246; Trichosteleum mastopomatoides, a new species from, 246
conduction, external water, strategy in bryophytes, 253+; technique for characterizing, 253+
Conocephalum conicum, 394+, 406
Cora pavonia, effects of desiccation on, 241
Corsinia, 252
Coxson, D. S. Effects of desiccation on net photosynthetic activity in the basidiomycete lichen Cora pavonia E. Fries from the cloud/mist zone of the tropical volcano La Soufrière (Guadeloupe), 241
Crandall-Stotler, B. *See* Stotler, R. E. & B. Crandall-Stotler, 50, 174, 265, 450
Crandall-Stotler, B., R. Stotler & P. Geissler. A biosystematic study of the subspecies of Fruillania tamarisci (L.), 287
Crocynia lesdainii (syn.), *163*; vouauxii (syn.), *163*
Crosby, M. R. Recent literature on mosses—136, 70; —137, 189; —138, 273
Crossomitrium patrisiae, 247
Crum, H. Bestia, Tripterocladium, and Isothecium: an explication of relationships, 40
crust soil associations, some lichens of, 48; Acarospora nodulosa var. nodulosa in, 48
Ctenidium malacodes, 249
Cyathodium barode, 251
Cyphelium tigillare, 213
cytology, of Mylia spp., 376+; of Pellia spp., 376+; of Plagiochila, 376+; of Riccia, 327

Dawsonia, 252
Dendroceros japonicus, 406+
Dermatocarpon acarosporoides (syn.), *28*; arboreum (syn.), *36*; compactum (syn.), *32*; cinereum (syn.), *31*; heppioides (syn.), *33*; lachneum (syn.), *35*; michelii (syn.), *35*; miniatum, 28; plumbeum, 36; rufescens (syn.), *34*; squamellum (syn.), *36*; trapeziforme (syn.), *35*; tuckermanii (syn.), *36*; weberi, 28; zahlbruckneri (syn.), *38*

desiccation, blue-green algae response to, 241+; effects on abscisic acid content in Anthoceros, 393+; effects on photosynthesis and distribution of Cora pavonia, 241; green algae response to, 241+
DEY, J. P. The distribution of the lichen, Bulbothrix isidiza, in North America north of Mexico, 417
Dicranella cerviculata, 43+; form, associated with coal deposits, 43+
Dicranolejeunea aberrans, 339; axillaris, 341; phyllorhiza, 339
Didymodon johansenii, 17
Diplophyllum imbricatum, 345+, mapped, 346
distribution, spatial, of moss colonies correlated with water relations, 254
Ditremis **limitans**, comb. nov., 163; **terminata**, comb. nov., 163
Dozya japonica, 409+, mapped, 414, peristome affinity to Leucodontaceae (peristome), 411*+; japonica var. robusta, 414
Dozyoideae subfam. Leucodontaceae, stat. nov., 413
DUCKETT, J. G. *See* RENZAGLIA, K. S. & J. G. DUCKETT, 419
Dumortiera dilatata, 372; hirsuta, 251+, 372

Ectropothecium leptochaeton, 247
EGAN, R. S. A fifth checklist of the lichen-forming, lichenicolous and allied fungi of the continental United States and Canada, 77
EGAN, R. S. Recent literature on lichens—129, 60; —130, 183; —131, 456
endemic Hepaticae in western North America, 344+
Endocarpon arboreum (syn.), *36*+; cinereum (syn.), *31*; exiguum (syn.), *35*; lachneus (syn.), *34*; michelii (syn.), *35*; pusillum var. rufescens (syn.), *34*; rufescens (syn.), *34*; tephroides (syn.), *31*; tuckermanii (syn.), *36*
Endopyrenium acarosporoides (syn.), *28*; americanum (syn.), *32*; bajadanae (syn.), *28*; crustaceum (syn.), *33*; granulosum (syn.), *33*; hepaticum (syn.), *34*; michelii (syn.), *35*; plumbeum (syn.), *36*; rufescens (syn.), *34*; rupicola (syn.), *32*; zahlbruckneri (syn.), *38*
endosporous germination, 405+
Enterographa **carnea**, comb. nov., 163
Entodon beyrichii, 247
epiphytic lichens, on trees in Yellowstone National Park, 212
Erythrodontium squarrosum, 247
euchromatin, in bryophytes, 376+
Eulacophyllum cultelliforme, 247
Eurhynchium semiscabrum, 247
EVERSMAN, S., C. JOHNSON & D. GUSTAFSON. Vertical distribution of epiphytic lichens on three tree species in Yellowstone National Park, 212
Exormotheca, abscisic acid in, 393+; bulbosa, 394+; holstii, 394
exosporous germination, 405+
exsiccati, Tierra del Fuego, species in, series I, nrs. 1–25, 76; series I, nrs. 26–75, 281.

Fissidens donnellii, 247; microcladus var. microcladus, 247; prionodes f. flexinervis, 247; prionodes f. hornschuchii, 247; prionodes f. puiggarii, 247; sharpii, 247; weirii var. hemicraspedophyllus, 247
flavonoid profiles, in Frullania tamarisci subspecies, 287+
fossil, of Stictolejeunea cf. squamata, 339
Fossombronia foveolata, 401+; hispidissima, 345+, mapped, 346; longiseta, 345+, mapped, 346
Frullania, species on Lord Howe Island, 365; aeolatis, 365+; asagrayana, 287+, 290*, 292*, 298*, 300*, 302*; californica, 346+, mapped, 346; chilcootiensis, 346+, mapped, 346; crawfordii, 370; densiloba, 406; ericoides, 365+; falciloba, 365+; falsa, 370; filipendra, 370; franciscana, 346+, mapped, 347; **fulfordiae**, sp. nov., 365, 366*; hattoriana, 346+, mapped, 347; howeana, 365+; moniliata, 289*+, 288*, 290*, 298*, 300*; moniliata ssp. breviramea, 289; moniliata ssp. obscura, 289; nisquallensis, 291+, 292*, 298*, 302*; obtusifolia, 370; pentapleura, 370; seriata, 370; squarrosa (syn.), *365*; **subincumbens**, sp. nov., 367, 368*; tamarisci, 288*, 292*, 300*; tamarisci ssp. asagrayana (syn.), *287*+; tamarisci ssp. moniliata (syn.), *287*+; tamarisci ssp. nisquallensis (syn.), *287*+, 346; tamarisci ssp. nisquallensis, mapped, 347; tamarisci ssp. obscura (syn.), *287*+; tamarisci ssp. obscura var. breviramea (syn.), *289*+; tamarisci ssp. tamarisci (syn.), *287*+; **vittiana**, sp. nov., 368, 369*; wildii, 370; yorkiana, 367
Frullanoides bahamensis, 339; corticalis, 339; densifolia, 339+; densifolia ssp. grandidentata, 338; laciniatiflora, 338; mexicana, 338; tristis, 339, mapped, 340
Fulford, Margaret H., a tribute to, 285
fungi, allied, lichen-forming, lichenocolous, checklist for the United States and Canada, 77+
Fuscidea **placidensis**, comb. nov., 163; **subfilamentosa**, comb. nov., 163
fusicoccin, effect on stomata of Anthoceros spp., 393+

Geisleria synchnogonioides (syn.), *164*
GEISSLER, P. *See* CRANDALL-STOTLER, B., R. STOTLER & P. GEISSLER, 287
Geothallus tuberosus, 347+, mapped, 347
β-glucosyl Yariv reagent, in detection of arabinogalactan protein, 401+
GRADSTEIN, S. R. The Ptychanthoideae of Latin America: an overview (studies on Lejeuneaceae subfamily Ptychanthoideae XVI), 337
GRIFFIN, D., III (review). J. Florschutz-De Waard (ed.), Flora of Suriname, Vol. VI, Part I, Musci (Part II), 198
Groutiella mucronifolia, 247
growth forms and water relations in bryophytes, 254
GUSTAFSON, D. *See* EVERSMAN, S., C. JOHNSON & D. GUSTAFSON, 212
Gymnocolea inflata, 401+
Gymnomitrion laceratum, 362, mapped, 363
Gyrothyra underwoodiana, 347+, mapped, 347

H^+, effects on Sphagnum species, 221
Haplomitrium, aligned with the Metzgeriales, 383+; blumei, 383+; gibbsiae, 383+; hookeri, 383+; mnioides, apical organization and leaf ontogeny in, 383+, 384*, 386, 388*, 401+
HARTUNG, W., E. W. WEILER & O. H. VOLK. Immunochemical evidence that abscisic acid is produced by several species of Anthocerotae and Marchantiales, 393
HATTORI, S. The Frullania flora of Lord Howe Island, 365
Helicodontium capillare, 247
Helicophyllum torquatum, 247
hepatics, abscisic acid in, 393+; arabinogalactan proteins in, 401+; developmental aspects of chromatin condensation in, 376+; distribution, of Lejeuneaceae subfam. Ptychanthoideae, 337+, of North American endemics, 345+, of noteworthy species, 362+, of species in the North Pacific Arc, 345+; ecological correlations of spore germination in, 405+; spermatogenesis in, 419; spore output in, 251+
Herbertus aduncus, 238; angustivittata, 374; sakuraii ssp. arcticus, 347+, mapped, 347
heterochromatin, in bryophytes, 376+
Heterodermia **erinacea**, comb. nov., 163

Heterophyllium affine, 247
HONG, W. S. The distribution of western North American Hepaticae. Endemic taxa and taxa with a North Pacific Arc distribution, 344
Hookeriopsis, sect. Hookeriopsis, 45+; **crosbyi**, sp. nov., 45, 46*; falcata, 247; websteri, new to Panama, 47
Hubbsia **californica**, comb. nov., 163; lumbricoides (syn.), *163*
hydroxyproline-containing proteins, in hepatics, 401
Hygrolejeunea alaskana, 344
Hygrophyla dilatata, 372
Hypnella pilifera, 247
Hypodontium, 231+
Hypogymnia austerodes, 213+; bitteri, 213

India, Syrrhopodon tjibodensis new to, 208
Isothecium, relationships in, 40+; alopecuroides, 42; brachycladon, 42; cristatum, 42; eumyosuroides, 42; myosuroides, 41+; stoloniferum, 41; viviparum, 42

Jaegerina scariosa, 247
Japan, Marchantias described by Stephani of, 331
JOHNSON, C. *See* EVERSMAN, S., C. JOHNSON & D. GUSTAFSON, 212
JOVET-AST, S. Vers une classification phylogénétique des espèces du genre Riccia, 321
Jubula japonica, 406
Julella **fallaciosa**, comb. nov., 163; **sublactea**, comb. nov., 163
Jungermannia exsertifolia ssp. cordifolia var. pendletonii, 347+, mapped, 347; diffusa (syn.), *374*; juliformis (syn.), *342*; leiantha, 401+; lunulata (syn.), *375*; rubra, 348+, mapped, 348; schusterana, 347+

KAUL, A. *See* PATIDAR, K. C., C. M. SOLANKI & A. KAUL, 251
KEENER, C. S. (review). W. F. Grant (ed.), Plant Biosystematics, 75
KOIDE, R. (review). D. C. Smith & A. E. Douglas, The Biology of Symbiosis, 197

lamellar strip, function in spermatogenesis, 441+
Latin America, Lejeuneaceae subfam. Ptychanthoideae in, 337+; Monoclea species in, 371+
Lecanora annulata (syn.), *163*; caesiopruinosa (syn.), *163*; limitata (syn.), *163*; mastoidea (syn.), *163*; narssaquensis (syn.), *163*; pertusa (syn.), *163*; piniperda, 212+; piniperda f. nigrescens, 213
Lecidea chlorosticta (syn.), *164*; granulata (syn.), *163*; placidensis (syn.), *163*
Lecidella **granulata**, comb. nov., 163
Lejeunea alaskana, 348+, mapped, 348; fuscescens (syn.), *342*; vaginata, 406
Lejeuneaceae subfam. Ptychanthoideae, in Latin America, 337+
Lembophyllaceae, genera of, 41
Lepidopilum polytrichoides, 247; scabrisetum, 247; tortifolium, 247
Lepidozia ceratophylla, 362
Lepraria **lesdainii**, comb. nov., 163; **vouauxii**, comb. nov., 163
Leskeodon andicola, 247+
Letharia columbiana, 213; vulpina, 213
Leucodon atrovireus (peristome), 412*, 413*; sapporensis (peristome), 413*
Leucodontaceae, peristome affinity of Dozya japonica, 409+; subfamily Leucodontoideae tribe Dozyeae (syn.), *413*
Leucodontopsis geniculata, 247
Leucomium lignicola, 247
Lichen chlorophyllus (syn.), *164*; erinaceus (syn.), *163*; juniperinus (syn.), *164*; lachneus (syn.), *34*; pinastri, 164; sepincola (syn.), *164*; tephroides (syn.), *31*
Lindigia aciculata, 247
Lopholejeunea howei, 339; muelleriana, 339; subfusca, 339
Lophozia holmenianum, 348+, mapped, 348
Lord Howe Island, Frullania species of, 365
Luteolejeunea herzogii, 338

Macromitrium, Serpotortella compared with, 232
MANKIEWICZ, P. S. The low pressure field porometer: a new, low cost technique for characterizing external capillary water conduction in whole colonies of bryophytes and other small plants, 253
Marchantia, typification and synonymy of Japanese spp., 331; alatocapitata, 331; calcarata, 331; crenata, 335+; cuneiloba (syn.), *331+*, 332*; fauriana (syn.), *333*; nepalensis, 251; paleacea ssp. diptera, 331+; paleacea ssp. paleacea, 333+; palmata, 251; pinnata, 333+, 334*; pinnatim-appendiculata (syn.), *333*; pinnatim-articulata (syn.), *333*; planipora, 333; polymorpha, 333+, 394; pulcherrima, 333; punaluana (syn.), *333*; tosana, 331+, 332*, 335*; tosayamensis (syn.), *335*; vaginata (syn.), *335+*
Marchantiales, abscisic acid in, 393+; spore output in, 251
Marchesinia brachiata, 341; robusta, 339
Marsupella bolanderi, 348+, mapped, 348
Mastigolejeunea auriculata, 339
MCCUNE, B. Distribution of chemotypes of Rhizoplaca in North America, 6
Melanelia **fuliginosa**, comb. nov., 163; **glabra**, comb. nov., 163; **granulosa**, comb. nov., 163; **incolorata**, comb. nov., 163
Melophyllum radiculosum, 309
Mesonodon flavescens, 247
Mesopyrenia sanfordensis (syn.), *163*
Micarea **chlorosticta**, comb. nov., 164; **endocyanea**, comb. nov., 164
Microcampylopus curvisetus, 247
mitochondria, in bryophyte spermatogenesis, 444+
Mittenothamnium reptans, 249
Mitthyridium, new synonyms and range records for paleotropic species of, 201+; binsteadii (syn.), *207*; chenagonii (syn.), *234*; cyrtophyllum (syn.), *232*; flavum, 207; letestui (syn.), *207*; parvulum (syn.), *207*; pungens (syn.), *207*; repens, 234; **retusum**, comb. nov., 201, 204*; subflavum (syn.), *208*; subluteum, 207; undulatum, new synonyms of, 207
Monoclea, distribution of, 371+; forsteri, 371+; gottschei (syn.), *371+*
MURRAY, B. M. Andreaea **schofieldiana** and A. **megistospora**, species novae, and taxonomic criteria for sect. Nerviae (Andreaeopsida), 15
Mycoporellum californicum (syn.), *164*
Mycoporum **compositum**, comb. nov., 164; sparsellum (syn.), *164*
Mylia, chromatin condensation in, 376+; anomala, 376+; taylorii, 376+

NEHIRA, K. Some ecological correlations of spore germination patterns in liverworts, 405
Nephroma americanum (syn.), *164*
Neurolejeunea catenulata, 339; seminervis, 339
New Zealand, Monoclea species in, 371+
NEWTON, M. E. Developmental aspects of chromatin condensation in liverworts, 376

NH_4^+, effects on Sphagnum species, 221
NO_3^-, effects on Sphagnum species, 221
North America, Acarospora nodulosa var. nodulosa new to, 48; Bulbothrix isidiza distribution north of Mexico in, 417+
North Pacific Arc, western North American Hepaticae of, 344+
Nowellia curvifolia, 401+
nuclear development, in spermatogenesis, 442+

Ocellularia **obturascens**, comb. nov., 164
Odontolejeunea lunulata, 339, 375
Odontoschisma gibbsiae, 348+, mapped, 349+; macounii, 237+; prostratum, 263
ontogeny, leaf, in Haplomitrium mnioides, 383
Orthostichidium guyanense, 248
Orthostichopsis crinita, 249
Oxymitra, 252; paleacea, 394+
Oxystegus tenuirostris 248+

paleotropics, new synonyms and range records of Calymperes, Mitthyridium and Syrrhopodon species in, 201+
Panama, Hookeriopsis crosbyi, a new species from, 45+; Hookeriopsis websteri new to, 47
parastomes, in Dozya japonica, 411
Parmelia exasperatula, 212+; fuliginosa f. incolorata (syn.), *164*; granulosa (syn.), *163*; olivacea α corticola a. glabra (syn.), *163*; olivacea γ fuliginosa (syn.), *163*; praesignis, 242; sulcata, 213+
Parmeliopsis ambigua, 213+
Parmentaria **nana**, comb. nov., 164
PATIDAR, K. C., C. M. SOLANKI & A. KAUL. Spore output in Marchantiales, 251
peatlands, Appalachian Mountain, acid precipitation a threat to, 221
Pellia, chromatin condensation in, 376+; borealis, 377; endiviaefolia, 406; epiphylla, 376+, phototropism in, 390+; neesiana, 377+
Peltigera polydactyla, 242; praetextata, 242
peristome, of Dozya japonica, 409+; of Leucodon, 412+
phenocopies, phyletic, induced in hepatics, 401+
phosphoglucoisomerase zymogram variation, in sspp. of Frullania tamarisci, 287+
photography, time-lapse, of phototropism in Pellia epiphylla, 390
Phragmicoma juliformis (syn.), *342*
phyletic phenocopies, induced in hepatics, 401+
Phyllogonium fulgens, 249
Phyllopsora subcorallina (syn.), *163*; subfilamentosa (syn.), *163*
phylogenetic classification, of Riccia, 321
Physcia adscendens, 213+
piedmont of the Appalachian Mountains, Syrrhopodon prolifer var. papillosus in, 263
Pilotrichella flexilis, 249
Pinus albicaulis, epiphytic lichens on, 212+; contorta, epiphytic lichens on, 212+
Pirrella cavifolia, 249
Placidiopsis, key to North American species, 38; cervinula, new to North America, 27+; custnani, 38; grappae, 38; minor, 27+
Placidium compactum (syn.), *32*; hepaticum var. pruinosum (syn.), *35*; hepaticum var. reticulatum (syn.), *35*; michelii (syn.), *35*
Placolecanora sphaeroidea, 11
Plagiocarpa **illota**, comb. nov., 164
Plagiochasma, abscisic acid in, 393+; appendiculatum, 251; intermedium, 251; rupestre, 394+
Plagiochila, chromatin condensation in, 376+; asplenioides, 378+; corniculata, 362, mapped, 363; porelloides, 378+; schofieldiana, 348+, mapped, 349; tridenticulata, 362
Plagiomnium rhynchophorum, 249
Plagiothecium schraderi, 248
plastids, in bryophyte spermatogenesis, 443+
Platyhypnidium aquaticum, 248
Platysma fendleri (syn.), *164*
Pleuridiopsis amblyocalyx (syn.), *311*
Pleuridium exsertum (syn.), *309*+; kieneri (syn.), *311*
Pleurotheliopsis nana (syn.), *164*
Pleurozia acinosa, 406
Pogonatum andinum, 248
Polyblastia fallaciosa (syn.), *163*
Polymeridium **pleiomerellum**, comb. nov., 164
Polytrichastrum, 4; xanthopilum (syn.), *1*
Polytrichum **appalachianum**, sp. nov., 1, 2*, cytology, 4, ecology, 4+, mapped, 4; commune, 1+; formosum, 1+; longisetum, 1+; ohioense, 3+; pallidisetum, 1+; xanthopilum (syn.), *1*+
Porella bolanderi, 348+, mapped, 349; navicularis, 348+; roellii, 348+, mapped, 349; roellii f. crispata, 348+, mapped, 349; roellii f. schofieldiana, 349+, mapped, 349
Porina wilsonii (syn.), *164*
porometer, low pressure, new, 253+
Porothamnium fasciculatum, 248
Porotrichum **vancouveriense**, comb. nov., 41
Pottiaceae, Calymperaceae and Serpotortellaceae compared with, 232
Prionodon densus, 249; fusco-lutescens, 250
protonema, aid in distribution of Dicranella cerviculata form, 43
Pseudocryphaea domingensis, 248
Ptychanthoideae subfam. of Lejeuneaceae in Latin America, 337+; Species present in Latin America, 341
Ptychanthus striatus, 406
Pyrenocollema **caesia**, comb. nov., 164; **halodytes**, comb. nov., 164; **strontianensis**, comb. nov., 164
Pyrenula **cruentata**, comb. nov., 164; **maculata**, stat. et comb. nov., 164; megalospora (syn.), *163*; nitidella var. maculata (syn.), *164*; **subelliptica**, comb. nov., 164

Racopilum tomentosum, 250
Radula bolanderi, 349+, mapped, 349; constricta, 406; obtusiloba ssp. polyclada, 349+, mapped, 350
Ramalina siliquosa, 7
Reboulia hemisphaerica, 251
REED, C. F. *See* ROBINSON, H. & C. F. REED, 43
REESE, W. D. Nomenclature of paleotropical Calymperaceae, with description of Syrrhopodon **meijeri**, sp. nov., 201
REESE, W. D. & R. H. ZANDER. Serpotortellaceae, a new family of mosses (Musci) from Madagascar and Reunion, 230
refugium, southern piedmont of the Appalachian Mountains, 263
rehydration recovery, of Cora pavonia, 241
Reinkella californica (syn.), *163*
Relicina abstrusa, 417; eximbricata, 417
RENZAGLIA, K. S. & J. G. DUCKETT. Spermatogenesis in Blasia pusilla: from young antheridial initial through mature spermatozoid, 419
Rhaphidostegium chlorocormum, 248
Rhizoplaca, key to North American species of, 13; North American distribution of chemotypes in, 6; baranowii, 10; chrysoleuca, 6+, chemical races in, 7, races mapped, 8; glaucophana, 7+, mapped, 9; haydenii, 7+, mapped, 9; marginalis, 7+, mapped, 9; melanophthalma, 6+,

chemical races of, 7, races mapped, 10; peltata, 6+, chemical races mapped, 11; subdiscrepans, 6
Rhodobryum beyrichianum, 248
Rhodocarpon compactum (syn.), *32*
Rhynchostegiopsis tunguraguana, 248
Riccardia sinuata, 401+
Riccia, 321+; chemistry of, 327; cytology of, 327; phylogenetic classification of, 321+; sect. Pilifer, 321+; albovestitata, 394+; angolensis, 394+; billardieri, 251; campbelliana, 349+, mapped, 350; caroliniana, 321+, 326*; crustata (spore), 325*; crystallina, 252; discolor, 251; duplex, 401+; duthiea, 394+; fluitans, 251+, (spore), 325*; frostii (spore), 325*; gangetica, 251; glauca, 252, 394+; gougetiana, 252, 394+; huebeneriana, 251; membranacea, 321+, (spore), 325*; perssonii, 321+, (spore), 325*; sommieri, 326*; sorocarpa (spore), 325*; stenophylla, 252; vitalii (spore), 325*
Rinodina marysvillensis var. thujae (syn.), *164*; **thujae**, comb. nov., 164
ROBINSON, H. & C. F. REED. Notes on the distribution of the Bartleya expression of Dicranella cerviculata in Appalachian coal fields, 43

Sagedia cinerea (syn.), *31*
SASTRE-DE JESÚS, I. *See* CHURCHILL, S. P. & I. SASTRE-DE JESÚS, 246
Scapania americana, 349+, mapped, 350; fulfordiae, 350+, mapped, 350, hians ssp. salishensis, 350+, mapped, 350; hollandiae, 350, mapped, 350; nemorosa, 401+; spitzbergensis, 237+; subalpina var. muddiae, 350, mapped, 350
Schiffneriolejeunea amazonica, 339; polycarpa, 339, mapped, 340
Schisma angustivittatum (syn.), 374
Schizomitrium rivulare, 248
Schofieldia monticola, 347+, mapped, 350
Schradeobryum ulicinum, 248
SCHUMACKER, R. *See* BISANG, I. & R. SCHUMACKER, 237
Sematophyllum **chlorocormum**, comb. nov., 248; constrictum, 248; oedophysidium, 248; subsimplex, 248; tequendamense, 248; turgidulum, 248
Serpotortella, 231+; key to species of, 232; Macromitrium compared with, 232; **chenagonii**, 230+, comb. nov., 234, 235*; **cyrtophylla**, 230+, comb. nov., 232, 233*; madagassa (syn.), *234*+; marginata (syn.), *234*+
Serpotortellaceae, fam. nov., 230+; Calymperaceae and Pottiaceae compared with, 232
sesquiterpene lactone profiles of Frullania tamarisci subspp., 304+
SHARP, A. J. Some hepatics with interesting distributions, 362
SNIDER, J. A. A revision of the moss genus Astomiopsis (Ditrichaceae), 309
SO_4^{2-}, effects on Sphagnum species, 221
SOLANKI, C. M. *See* PATIDAR, K. C., C. M. SOLANKI & A. KAUL, 251
spermatogenesis, in Blasia pusilla, 430+
spermatogenic processes, systematic considerations of, 445
Sphagnum, hollow-dwelling species response to acidity, 224+; hummock-dwelling species response to acidity, 224+; affine, 217+; austinii, 217+; carlottae (syn.), *217*; fallax, effects of acidity on, 221+; henryense, effects of acidity on, 221+; imbricatum, 217+; imbricatum complex, key to, 218, table of characters, 219; imbricatum ssp. affine (syn.), *217*+; imbricatum ssp. affine var. flagellare (syn.), *217*+; imbricatum ssp. austinii (syn.), *217*+; imbricatum ssp. austinii var. arcticum (syn.), *218*; portoricense, 217+; pulchrum, effects of acidity on, 221+; steerei, 217+, stat. nov., 218
spore, germination patterns in liverworts, 405+; morphology in Riccia, phylogenetic interpretation of, 321; output in Marchantiales, 251+; spore size in systematics of Andreaeae, sect. Nerviae, 25
sporeling types, hepatic, correlated with habitat, 405+
Spruceanthus theobromae, 338
Squamidium leucotrichum, 250
ST. CLAIR, L. L. & R. B. WARRICK. Acarospora nodulosa (Duf.) Hue. v. nodulosa: a new record for North America, 48
Steereobron subulirostrum, 248
Stephani, Japanese Marchantia species described by, 331
Stereocaulon virgatum, 241+
Stereophyllum radiculosum, 248
Stictolejeunea balfourii, 339+, mapped, 340; balfourii var. bekkeri, 338
stomata, chemical mechanism in Anthoceros, 393+
STOTLER, R. *See* CRANDALL-STOTLER, B., R. STOTLER & P. GEISSLER, 287
STOTLER, R. E. Margaret H. Fulford—a tribute, 285
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—109, 50; —110, 174; —111, 265; —112, 450
Strigula **submuriformis**, comb. nov., 164; **synchnogonioides**, comb. nov., 164; **wilsonii**, comb. nov., 164
Sumatra, Syrrhopodon tjibodensis new to, 208
Sutherland Falls, Blanket Creek Provincial Park, British Columbia, rare liverworts in, 237+
Symbiezidium transversale ssp. dentatum, 338
Syrrhopodon, new synonyms and range records for paleotropic species of, 201+; armatus, 208; armatus var. latifolius (syn.), *208*; assamicus (syn.), *208*; binsteadii (syn.), *207*; burmensis (syn.), *206*; chenagonii (syn.), *230*+; crispus, excluded from Calymperaceae, 208; cryptocarpos (syn.), *248*; cymbifolius (syn.), *248*; cyrtophyllus (syn.), *230*+; gaudichaudii, 208; heterophyllus (syn.), *206*; himalayanus (syn.), *208*; hornschuchii, 248; incompletus, 263+; incompletus var. incompletus, 208, African report of, 208; jacobii (syn.), *234*+; leptodontoides (syn.), *232*+; linealis (syn.), *208*; **meijeri**, sp. nov., 201, 202*; nossibeanus (syn.), *207*; parsiticus, 248, 263+; parasiticus var. disciformis, 208; parasiticus var. parasiticus, 208; parvulus (syn.), *207*; prolifer var. papillosus, new to Alabama, 263; prolifer var. prolifer in Alabama, 263; pungens (syn.), *207*; robustiusculus, new synonyms of, 208; **sarawakensis**, comb. nov., 203; subflavus (syn.), *208*; texanus, 263; tjibodensis, new to India, Thailand and Sumatra, 208; tortelloides (syn.), *234*; tristichus, 208

Takakia, 388; ceratophylla, 362, mapped, 363; lepidozioides, 362, mapped, 363
TAN, B. C. (review). Z. Iwatsuki & M. Izawa, Shida, Koke [Pteridophytes and Bryophytes]. Nature Handbook 13, 280
Targionia hypophylla, 251
TAXACOM, electronic bulletin board system, 76
terpenoid profiles, in Frullania tamarisci subspecies, 287+
Thailand, Syrrhopodon tjibodensis new to, 208
Thelotrema bahianum var. obturascens (syn.), *164*; carneum (syn.), *163*
THOMAS, R. J., P. J. CARON & R. S. WATT. Time-lapse photography of phototropism in Pellia epiphylla, 390
THOMSON, J. W. The lichen genera Catapyrenium and Placidiopsis in North America, 27
Thyridium binsteadii (syn.), *207*; chenagonii (syn.), *234*; cyrtophyllum (syn.), *232*; letestui (syn.), *207*; letestui var. scalariforme (syn.), *207*; parvulum (syn.), *207*; subflavum (syn.), *208*

Thysananthus pterobryoides, 339
Tierra del Fuego, species in Musci Fuegiani Exsiccati: series I, nrs. 1–25, 76; species in Musci Fuegiani Exsiccati: series I, nrs. 26–75, 281
TIMME, S. L. & D. W. WORLEY. Syrrhopodon prolifer Schwaegr. var. papillosus (C. Muell.) Reese in Alabama: a new state record, 263
Tomasellia **californica**, comb. nov., 164; **sparsella**, comb. nov., 164
Tortella tortuosa, 248
Trematodon longicollis, 248
Trichocoleopsis sacculata, 406
Trichosteleum mastopomatoides, 249*; **mastopomatoides**, sp. nov., 248; papillosum, 249; subdemissum, 249
Tripterocladium, placed in Hypnaceae, 40; relationships in, 40+; leucocladium, 41
Tuckermannopsis **americana**, comb. nov., 164; **aurescens**, comb. nov., 164; **canadensis**, comb. nov., 164; **chlorophylla**, comb. nov., 164; **fendleri**, comb. nov., 164; **juniperina**, comb. nov., 164; **merrillii**, comb. nov., 164; **oakesiana**, comb. nov., 164; **pallidula**, comb. nov., 164; **pinastri**, comb. nov., 164; **platyphylla**, comb. nov., 164; **sepincola**, comb. nov., 164; **viridis**, comb. nov., 164

Usnea fulvoreagens, 213+
Utah, Acarospora nodulosa var. nodulosa in, 48

Verdoornianthus griffinii, 338; marsupiifolius, 339
Verrucaria caesia (syn.), *164*; cervinula (syn.), *38*; distans (syn.), *163*; halodytes (syn.), *164*; illota (syn.), *164*; limitans (syn.), *163*; plumbaria (syn.), *163*; prasina (syn.), *163*; squamella (syn.), *36*; strontianensis (syn.), *164*; subelliptica (syn.), *164*; sublactea (syn.), *163*; terminata (syn.), *163*; viridiseda f. albiseda (syn.), *163*
VOLK, O. H. *See* HARTUNG, W., E. W. WEILER & O. H. VOLK, 393

WARRICK, R. B. *See* ST. CLAIR, L. L. & R. B. WARRICK, 48
WATT, R. S. *See* THOMAS, R. J., P. J. CARON & R. S. WATT, 390
WEILER, E. W. *See* HARTUNG, W., E. W. WEILER & O. H. VOLK, 393
WIEDER, R. K. *See* AUSTIN, K. A. & R. K. WIEDER, 221
WORLEY, D. W. *See* TIMME, S. L. & D. W. WORLEY, 263

Xanthoria fallax, 213+

YANO, O. Additions to the Brazilian Hepaticae, 374
Yellowstone National Park, epiphytic lichens in, 212

ZANDER, R. H. *See* REESE, W. D. & R. H. ZANDER, 230
Zoopsis antillana, new to Brazil, 374